

ENHANCED SPECIALTY GUIDELINE MANAGEMENT

KALBITOR (ecallantide)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication

Treatment of acute attacks of hereditary angioedema (HAE) in patients 12 years of age and older

All other indications are considered experimental/investigational and are not a covered benefit.

II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

- A. C4 levels and C1 inhibitor functional and antigenic protein levels
- B. F12, angiotensin-converting enzyme 1 or plasminogen gene mutation testing, if applicable
- C. Chart notes confirming family history of angioedema, if applicable

III. CRITERIA FOR INITIAL APPROVAL

Authorization of 12 months may be granted for treatment of acute hereditary angioedema attacks when the requested medication will not be used in combination with Berinert, Firazyr, or Ruconest and either of the following criteria is met:

- A. Member has C1 inhibitor deficiency or dysfunction as confirmed by laboratory testing:
 1. C1 inhibitor (C1-INH) antigenic level is below the lower limit of normal as defined by the laboratory performing the test, or
 2. Normal C1-INH antigenic level and a low C1-INH functional level (functional C1-INH less than 50% or C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test).
- B. Member has normal C1 inhibitor as confirmed by laboratory testing and meets one of the following criteria:
 1. Member has an F12, angiotensin-converting enzyme 1, or plasminogen gene mutation as confirmed by genetic testing, or
 2. Member has a documented family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine (e.g., cetirizine) for at least one month.

IV. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for continuation of therapy when all of the following criteria are met:

- A. Member meets the criteria for initial approval.
- B. Member has experienced reduction in severity and/or duration of attacks when they use the requested medication to treat an acute attack.

V. REFERENCES

1. Kalbitor [package insert]. Burlington, MA: Dyax Corp.; March 2015.
2. Bowen T, Cicardi M, Farkas H, et al. 2010 International consensus algorithm for the diagnosis, therapy, and management of hereditary angioedema. *Allergy Asthma Clin Immunol*. 2010;6(1):24.
3. Cicardi M, Bork K, Caballero T, et al. Hereditary Angioedema International Working Group. Evidence-based recommendations for the therapeutic management of angioedema owing to hereditary C1 inhibitor deficiency: consensus report of an International Working Group. *Allergy*. 2012;67:147-157.
4. Zuraw BL, Banerji A, Bernstein JA, et al. US Hereditary Angioedema Association Medical Advisory Board 2013 recommendations for the management of hereditary angioedema due to C1 inhibitor deficiency. *J Allergy Clin Immunol: In Practice*. 2013; 1(5): 458-467.
5. Zuraw BL, Bork K, Binkley KE, et al. Hereditary angioedema with normal C1 inhibitor function: consensus of an international expert panel. *Allergy Asthma Proc*. 2012; 33(6):S145-S156.
6. Maurer M, Magerl M, Ansotegui I, et al. The international WAO/EAACI guideline for the management of hereditary angioedema – the 2017 revision and update. *Allergy*. 2018;00:1-22.
7. Lang DM, Aberer W, Bernstein JA, et al. International consensus on hereditary and acquired angioedema. *Ann Allergy Asthma Immunol*. 2012; 109:395-202.
8. Cicardi M, Aberer W, Banerji A, et al. Classification, diagnosis, and approach to treatment for angioedema: consensus report from the Hereditary Angioedema International Working Group. *Allergy*. 2014;69: 602-616.
9. Bowen T. Hereditary angioedema: beyond international consensus – circa December 2010 – The Canadian Society of Allergy and Clinical Immunology Dr. David McCourtie Lecture. *Allergy Asthma Clin Immunol*. 2011;7(1):1.
10. Bernstein J. Update on angioedema: Evaluation, diagnosis, and treatment. *Allergy and Asthma Proceedings*. 2011;32(6):408-412.
11. Longhurst H, Cicardi M. Hereditary angio-edema. *Lancet*. 2012;379:474-481.
12. Farkas H, Martinez-Saguer I, Bork K, et al. International consensus on the diagnosis and management of pediatric patients with hereditary angioedema with C1 inhibitor deficiency. *Allergy*. 2017;72(2):300-313.
13. Henao MP, Kraschnewski J, Kelbel T, Craig T. Diagnosis and screening of patients with hereditary angioedema in primary care. *Therapeutics and Clin Risk Management*. 2016; 12: 701-711.