

Reference number(s)
2516-A

SPECIALTY GUIDELINE MANAGEMENT

SYMDEKO (tezacaftor/ivacaftor)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication¹

Symdeko is indicated for the treatment of patients with cystic fibrosis (CF) aged 6 years and older who are homozygous for the *F508del* mutation or who have at least one mutation in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene that is responsive to tezacaftor/ivacaftor based on *in vitro* data and/or clinical evidence. If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of *CFTR* mutation followed by verification with bi-directional sequencing when recommended by the mutation test instructions for use.

All other indications are considered experimental/investigational and are not a covered benefit.

II. REQUIRED DOCUMENTATION

The following information is necessary to initiate the prior authorization review: genetic testing report confirming the presence of the appropriate *CFTR* gene mutation.

III. CRITERIA FOR INITIAL APPROVAL

Cystic Fibrosis¹⁻³

Indefinite authorization may be granted for treatment of cystic fibrosis when all of the following criteria are met:

- A. Genetic testing was conducted to detect a mutation in the *CFTR* gene.
- B. The member has one of the following mutations in the *CFTR* gene: A455E, A1067T, D110E, D110H, D579G, D1152H, D1270N, E56K, E193K, E831X, F1052V, F1074L, K1060T, L206W, P67L, R74W, R117C, R347H, R352Q, R1070W, S945L, S977F, 711+3A→G, 2789+5G→A, 3272-26A→G, 3849+10kbC→T, or the member is homozygous for the *F508del* mutation.
- C. The member is at least 6 years of age.
- D. Symdeko will not be used in combination with Kalydeco or Orkambi.

IV. CONTINUATION OF THERAPY

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

V. REFERENCES

1. Symdeko [package insert]. Boston, MA: Vertex Pharmaceuticals Inc.; June 2019.

Symdeko_PA_ALL_Rx

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

Reference number(s)
2516-A

2. Rowe SM, Daines C, Ringshausen FC, Kerem E, Wilson J, Tullis E, Nair N, Simard C, Han L, Ingenito EP, McKee C, Lekstrom-Himes J, Davies JC. Tezacaftor-ivacaftor in Residual Function Heterozygotes with Cystic Fibrosis. *N Engl J Med*. 2017; 377:2024-2035
3. Taylor-Cousar JL, Munck A, McKone EF, et al. Tezacaftor-ivacaftor in patients with cystic fibrosis homozygous for Phe508del *N Engl J Med* 2017; 377:2013-2023