

**Taltz® (ixekizumab)  
Effective 02/20/19**

<b>Plan</b>	<input checked="" type="checkbox"/> MassHealth <input type="checkbox"/> Commercial/Exchange	<b>Program Type</b>	<input checked="" type="checkbox"/> Prior Authorization
<b>Benefit</b>	<input checked="" type="checkbox"/> Pharmacy Benefit <input type="checkbox"/> Medical Benefit (NLX)		<input type="checkbox"/> Quantity Limit <input type="checkbox"/> Step Therapy
<b>Specialty Limitations</b>	This medication has been designated specialty and must be filled at a contracted specialty pharmacy.		
<b>Contact Information</b>	<b>Specialty Medications</b>		
	All Plans	Phone: 866-814-5506	Fax: 866-249-6155
	<b>Non-Specialty Medications</b>		
	MassHealth	Phone: 877-433-7643	Fax: 866-255-7569
	Commercial	Phone: 800-294-5979	Fax: 888-836-0730
	Exchange	Phone: 855-582-2022	Fax: 855-245-2134
	<b>Medical Specialty Medications (NLX)</b>		
	All Plans	Phone: 844-345-2803	Fax: 844-851-0882
<b>Exceptions</b>	N/A		

**Overview**

Taltz® (ixekizumab) is a humanized interleukin-17A antagonist indicated for the treatment of adults with:

1. Moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy
2. Active psoriatic arthritis
3. Active ankylosing spondylitis

**Coverage Guidelines**

Approval will be granted if the member meets the following diagnosis specific criteria:

**Moderate to severe plaque psoriasis**

1. Member has a diagnosis of moderate-severe plaque psoriasis **AND**
2. Member is at least 18 years of age **AND**
3. Prescriber has provided documentation of ONE of the following:
  - a. Inadequate response, adverse reaction, or contraindication to at least TWO conventional therapies in any one of the following combinations (combinations DO NOT have to be used concurrently):
    - i. 1 topical agent + 1 systemic agent
    - ii. 1 topical agent + 1 phototherapy (required for diagnosis of guttate psoriasis)
    - iii. 1 systemic agent + 1 phototherapy
    - iv. 2 systemic agents
  - b. Contraindication to ALL conventional therapies
    - i. Topical agents
    - ii. Phototherapy
    - iii. Systemic agents
  - c. Inadequate response or adverse reaction to ONE biologic DMARD that is FDA-approved



for plaque psoriasis

**AND**

4. Dosing is appropriate (see appendix)

**Psoriatic arthritis**

1. Member has a diagnosis of psoriatic arthritis **AND**
2. Member is at least 18 years of age **AND**
4. Prescriber has provided documentation of ONE of the following:
  - a. Inadequate response, adverse reaction, or contraindication to at least ONE traditional DMARD (hydroxychloroquine, methotrexate, sulfasalazine)
  - b. Inadequate response or adverse reaction to ONE biologic DMARD that is FDA-approved for the requested indication

**AND**

5. Dosing is appropriate (see appendix)

**Continuation of Therapy**

Reauthorization requires physician documentation indicating a positive response to therapy.

**Limitations**

1. Initial approvals will be varied based on the treatment:
  - a. For moderate to severe plaque psoriasis, approvals will be granted for 3 months.
  - b. For psoriatic arthritis, approvals will be granted for 6 months.
2. Reauthorizations will be granted for 12 months.

**Appendices**

**Dosing**

<p>Taltz® (ixekizumab)</p>	<p><b>Plaque Psoriasis:</b>  <u>SQ:</u> 160 mg initially at week 0, followed by 80 mg at weeks 2, 4, 6, 8, 10 and 12; followed by 80 mg every four weeks</p> <p><b>Psoriatic Arthritis:</b>  <u>SQ:</u> 160 mg initially at week 0, followed by 80 mg every 4 weeks.</p> <p>For psoriatic arthritis patients with coexistent moderate-to-severe plaque psoriasis, use the dosing regimen for plaque psoriasis</p>
--------------------------------	---

**References**

1. Taltz (ixekizumab) [prescribing information]. Indianapolis, IN: Eli Lilly and Co; May 2018
2. Nash P, Kirkham B, Okada M, et al. Ixekizumab for the treatment of patients with active psoriatic arthritis and an inadequate response to tumour necrosis factor inhibitors: results from the 24-week randomised, double-blind, placebo-controlled period of the SPIRIT-P2 phase 3 trial. Lancet 2017; 389:2317.
3. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. J Am Acad Dermatol. 2009; 61:451-485.



4. Griffiths CE, Reich K, Lebwohl M, et al. Comparison of ixekizumab with etanercept or placebo in moderate-to-severe psoriasis (UNCOVER-2 and UNCOVER-3): results from two phases 3 randomised trials. *Lancet*. 2015;386(9993):541-51.
5. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2016;75(3):499-510.
6. McInnes IB, Mease PJ, Kirkham B, et al. Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2015;386(9999):1137-46.

### **Review History**

11/28/16 – Reviewed

02/27/17 – Adopted SGM & ST

03/20/17 – Effective

03/01/18 – Adopted MH RS

02/20/19 – Reviewed in P&T Meeting

### **Disclaimer**

AllWays Health Partners complies with applicable federal civil rights laws and does not discriminate or exclude people on the basis of race, color, national origin, age, disability, or sex.