

<b>Reference number</b>
1734-A

## SPECIALTY GUIDELINE MANAGEMENT

### SANDOSTATIN (octreotide acetate injection) SANDOSTATIN LAR DEPOT (octreotide acetate for injectable suspension) octreotide acetate injection

#### POLICY

#### I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### A. FDA-Approved Indications

1. octreotide acetate/Sandostatin:
  - a. Indicated to reduce blood levels of growth hormone and IGF-1 (somatomedin C) in acromegaly patients who have had inadequate response to or cannot be treated with surgical resection, pituitary irradiation, and bromocriptine mesylate at maximally tolerated doses.
  - b. Indicated for the symptomatic treatment of patients with metastatic carcinoid tumors where it suppresses or inhibits the severe diarrhea and flushing episodes associated with the disease.
  - c. Indicated for the treatment of the profuse watery diarrhea associated with vasoactive intestinal peptide (VIP)-secreting tumors.
2. Sandostatin LAR: Sandostatin LAR Depot is indicated in patients in whom initial treatment with Sandostatin injection has been shown to be effective and tolerated.
  - a. Indicated for long-term maintenance therapy in acromegalic patients who have had an inadequate response to surgery and/or radiotherapy, or for whom surgery and/or radiotherapy is not an option.
  - b. Indicated for long-term treatment of the severe diarrhea and flushing episodes associated with metastatic carcinoid tumors.
  - c. Indicated for long-term treatment of the profuse watery diarrhea associated with vasoactive intestinal peptide (VIP)-secreting tumors.

##### B. Compendial Uses

1. Neuroendocrine tumors (NETs):
  - a. Tumors of the gastrointestinal (GI) tract, lung, and thymus (carcinoid tumors)
  - b. Tumors of the pancreas
2. Pheochromocytoma and paraganglioma
3. Meningiomas
4. Thymomas and thymic carcinomas
5. Congenital hyperinsulinism (CHI)/persistent hyperinsulinemic hypoglycemia of infancy (PHHI) (octreotide and Sandostatin only)
6. Acquired immune deficiency syndrome (AIDS)-associated diarrhea
7. Inoperable bowel obstruction
8. Chemotherapy- and radiation-induced diarrhea
9. Enterocutaneous fistula
10. Gastroesophageal varices
11. Islet cell tumors
12. Pancreatic fistulas
13. Pituitary adenoma

octreotide (Sandostatin), Sandostatin LAR\_PA\_ALL\_MBRx

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.

<b>Reference number</b>
1734-A

14. Short bowel syndrome
15. Zollinger-Ellison syndrome

All other indications are considered experimental/investigational and are not a covered benefit.

## II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

- A. For acromegaly:
  1. For initial approval: Laboratory report indicating high pretreatment insulin-like growth factor-1 (IGF-1) level and chart notes indicating an inadequate or partial response to surgery or radiotherapy or a clinical reason for not having surgery or radiotherapy.
  2. For continuation: Laboratory report indicating normal current IGF-1 levels or chart notes indicating that the member's IGF-1 level has decreased or normalized since initiation of therapy
- B. Chemotherapy- and radiation-induced diarrhea: Chart notes indicating grade 3 or 4 diarrhea with current chemotherapy or radiation.

## III. CRITERIA FOR INITIAL APPROVAL

### A. Acromegaly

Authorization of 12 months may be granted for the treatment of acromegaly when all of the following criteria are met:

1. Member has a high pretreatment IGF-1 level for age and/or gender based on the laboratory reference range.
2. Member had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason why the member has not had surgery or radiotherapy.

### B. Neuroendocrine tumors (NETs)

1. Tumors of the gastrointestinal (GI) tract (carcinoid tumor)  
Authorization of 12 months may be granted for treatment of locoregional advanced or metastatic NETs of the GI tract or unresected primary gastrinoma.
2. Tumors of the thymus (carcinoid tumor)  
Authorization of 12 months may be granted for treatment of unresectable or metastatic NETs of the thymus.
3. Tumors of the lung (carcinoid tumor)  
Authorization of 12 months may be granted for treatment of unresectable or metastatic NETs of the lung.
4. Tumors of the pancreas  
Authorization of 12 months may be granted for treatment of NETs of the pancreas.

### C. Carcinoid syndrome

Authorization of 12 months may be granted for treatment of carcinoid syndrome when it is used in any of the following clinical settings:

1. As a single agent
2. In combination with telotristat for persistent diarrhea due to poorly controlled carcinoid syndrome
3. In combination with other systemic therapy options for persistent symptoms such as flushing or diarrhea, or for progressive disease

<b>Reference number</b>
1734-A

**D. Vasoactive intestinal peptide tumors (VIPomas)**

Authorization of 12 months may be granted for management of symptoms related to hormone hypersecretion of VIPomas.

**E. Meningiomas**

Authorization of 12 months may be granted for treatment of unresectable recurrent or progressive meningioma.

**F. Pheochromocytoma and paraganglioma**

Authorization of 12 months may be granted for treatment of locally unresectable or metastatic pheochromocytoma and paraganglioma.

**G. Thymomas and thymic carcinomas**

Authorization of 12 months may be granted for treatment of thymomas and thymic carcinomas when the requested drug is used as a second-line therapy with or without prednisone in any of the following clinical settings:

1. Unresectable disease following first-line chemotherapy for potentially resectable locally advanced disease, solitary metastasis, or ipsilateral pleural metastasis
2. Extrathoracic metastatic disease

**H. Congenital hyperinsulinism (CHI)/persistent hyperinsulinemic hypoglycemia of infancy (octreotide and Sandostatin only)**

Authorization of 6 months may be granted for treatment of CHI and persistent hyperinsulinemic hypoglycemia in an infant.

**I. AIDS-associated diarrhea**

Authorization of 12 months may be granted for treatment of AIDS-associated severe secretory diarrhea when anti-microbial (e.g., ciprofloxacin or metronidazole) or anti-motility agents (e.g., loperamide or diphenoxylate and atropine) have become ineffective.

**J. Bowel obstruction in terminal cancer**

Authorization of 12 months may be granted for management of GI symptoms (e.g., nausea, pain, vomiting) of inoperable bowel obstruction in members with terminal cancer.

**K. Chemotherapy- and radiation-induced diarrhea**

Authorization of 12 months may be granted for treatment of chemotherapy- or radiation-induced diarrhea when any of the following criteria are met:

1. Member is receiving treatment with chemotherapy or radiation
2. Member has grade 3 or greater diarrhea according to National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events (CTCAE).

**L. Enterocutaneous fistula**

Authorization of 12 months may be granted for management of volume depletion from enterocutaneous fistula.

**M. Gastroesophageal varices**

Authorization of 6 months may be granted for treatment of acute bleeding of gastroesophageal varices associated with cirrhosis.

**N. Islet cell tumors**

Authorization of 12 months may be granted for stabilization of blood glucose levels in patients with functioning islet cell tumors (e.g., insulinomas or glucagonomas).

<b>Reference number</b>
1734-A

**O. Pancreatic fistulas**

Authorization of 6 months may be granted for prevention and treatment of pancreatic fistulas following pancreatic surgery.

**P. Pituitary adenoma**

Authorization of 12 months may be granted for treatment of pituitary adenoma.

**Q. Short bowel syndrome**

Authorization of 12 months may be granted for treatment of short bowel syndrome when the daily intravenous fluid requirement is greater than 3 liters.

**R. Zollinger-Ellison syndrome**

Authorization of 12 months may be granted for treatment of Zollinger-Ellison syndrome.

**IV. CONTINUATION OF THERAPY**

**A. Acromegaly**

Authorization of 12 months may be granted for continuation of therapy for acromegaly when the member's IGF-1 level has decreased or normalized since initiation of therapy.

**B. Carcinoid syndrome, VIPomas, AIDS-associated diarrhea, bowel obstruction, chemotherapy/radiation-induced diarrhea, islet cell tumors, and Zollinger-Ellison syndrome**

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization when the member is experiencing clinical benefit as evidenced by improvement or stabilization in clinical signs and symptoms since initiation of therapy.

**C. All other indications**

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

**V. REFERENCES**

1. Octreotide acetate [package insert]. Rockford, IL: Mylan Institutional LLC; September 2017.
2. Sandostatin [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; April 2019.
3. Sandostatin LAR Depot [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; April 2019.
4. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: <http://www.nccn.org>. Accessed June 11, 2019.
5. Katznelson L, Laws ER, Melmed S, et al. Acromegaly: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2014;99:3933-3951.
6. American Association of Clinical Endocrinologists Acromegaly Guidelines Task Force. Medical guidelines for clinical practice for the diagnosis and treatment of acromegaly – 2011 update. *Endocr Pract*. 2011;17(suppl 4):1-44.
7. The NCCN Clinical Practice Guidelines in Oncology® Neuroendocrine Tumors (Version 1.2019). © 2019 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed June 11, 2019.
8. Rinke A, Muller H, Schade-Brittinger C, et al. Placebo-controlled, double-blind, prospective, randomized study on the effect of octreotide LAR in the control of tumor growth in patients with metastatic neuroendocrine midgut tumors: a report from the PROMID study group. *J Clin Oncol*. 2009;27:4656-4663.

Reference number
1734-A

9. The NCCN Clinical Practice Guidelines in Oncology® Central Nervous System Cancers (Version 2.2018). © 2018 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed January 26, 2019.
10. The NCCN Clinical Practice Guidelines in Oncology® Thymomas and Thymic Carcinomas. (Version 2.2018). © 2018 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed January 26, 2019.
11. The NCCN Clinical Practice Guidelines in Oncology® Palliative Care (Version 2.2019). © 2019 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed June 11, 2019.
12. Harris AG, O'Dorisio TM, Woltering EA, et al. Consensus statement: Octreotide dose titration in secretory diarrhea. Diarrhea Management Consensus Development Panel. *Dig Dis Sci*. 1995;40(7):1464-1473.
13. Fried M. Octreotide in the treatment of refractory diarrhea. *Digestion*. 1999;60:42-46.
14. Edmunds MC, Chen JD, Soykan I, et al. Effect of octreotide on gastric and small bowel motility in patients with gastroparesis. *Aliment Pharmacol Ther*. 1998;12(2):167-174.
15. Ripamonti C, Mercadante S, Groff L, et al. Role of octreotide, scopolamine butylbromide, and hydration in symptom control of patients with inoperable bowel obstruction and nasogastric tubes: A prospective randomized trial. *J Pain Symptom Manage*. 2000;19(1):23-34.
16. Mercadante S, Ripamonti C, Casuccio A, et al. Comparison of octreotide and hyoscine butylbromide in controlling gastrointestinal symptoms due to malignant inoperable bowel obstruction. *Support Care Cancer*. 2000;8(3):188-191.
17. Dorta G. Role of octreotide and somatostatin in the treatment of intestinal fistulae. *Digestion*. 1999;60 Suppl 2:53-56.
18. Jamil M, Ahmed U, Sobia H. Role of somatostatin analogues in the management of enterocutaneous fistulae. *J Coll Physicians Surg Pak*. 2004;14(4):237-240.
19. Rahbour G, Siddiqui MR, Ullah MR, et al. A meta-analysis of outcomes following use of somatostatin and its analogues for the management of enterocutaneous fistulas. *Ann Surg*. 2012;256(6):946-954.
20. Freitas DS, Sofia C, Pontes JM, et al. Octreotide in acute bleeding esophageal varices: A prospective randomized study. *Hepatogastroenterology*. 2000;47(35):1310-1314.
21. Imperiale TF, Teran JC, McCullough AJ. A meta-analysis of somatostatin versus vasopressin in the management of acute esophageal variceal hemorrhage. *Gastroenterology*. 1995;109(4):1289-1294.
22. Gøtzsche PC, Hróbjartsson A. Somatostatin analogues for acute bleeding oesophageal varices. *Cochrane Database Syst Rev*. 2008;(3):CD000193.
23. Erstad BL. Octreotide for acute variceal bleeding. *Ann Pharmacother*. 2001;35(5):618-626.
24. Corley DA, Cello JP, Adkisson W, et al. Octreotide for acute esophageal variceal bleeding: A meta-analysis. *Gastroenterology*. 2001;120(4):946-954.
25. Gross M, Schiemann U, Muhlhofer A, Zoller WG. Meta-analysis: Efficacy of therapeutic regimens in ongoing variceal bleeding. *Endoscopy*. 2001;33(9):737-746.
26. Li-Ling J, Irving M. Somatostatin and octreotide in the prevention of postoperative pancreatic complications and the treatment of enterocutaneous pancreatic fistulas: A systematic review of randomized controlled trials. *Br J Surg*. 2001;88(2):190-199.
27. Machado NO. Pancreatic fistula after pancreatectomy: Definitions, risk factors, preventive measures, and management – Review. *Int J Surg Oncol*. 2012;2012:602478.
28. Adachi T, Kuroki T, Kitasato A, et al. Safety and efficacy of early drain removal and triple-drug therapy to prevent pancreatic fistula after distal pancreatectomy. *Pancreatol* 2015;15:411-416.
29. Gurusamy KS, Koti R, Fusai G, Davidson BR. Somatostatin analogues for pancreatic surgery. *Cochrane Database Syst Rev*. 2013;4:CD008370.
30. Alberta Provincial CNS Tumour Team. Pituitary adenomas. Clinical Practice Guideline No. CNS-006 Edmonton, AB: Alberta Health Services, Cancer Care; August 2012.
31. Peeters M, Van den Brande J, Francque S. Diarrhea and the rationale to use Sandostatin. *Acta Gastroenterol Belg*. 2010;73(1):25-36.
32. Mystakidou K, Tsilika E, Kalaidopoulou O, et al. Comparison of octreotide administration vs conservative treatment in the management of inoperable bowel obstruction in patients with far advanced cancer: A randomized, double-blind, controlled clinical trial. *Anticancer Res*. 2002;22(2B):1187-1192.

Reference number
1734-A

33. American Gastroenterological Association. American Gastroenterological Association medical position statement: Short bowel syndrome and intestinal transplantation. *Gastroenterology*. 2003;124(4):1105-1110.
34. Loehrer PJ Sr, Wang W, Johnson DH, et al. Octreotide alone or with prednisone in patients with advanced thymoma and thymic carcinoma: An Eastern Cooperative Oncology Group Phase II Trial. *J Clin Oncol*. 2004;22(2):293-299.
35. Leandros E, Antonakis PT, Albanopoulos K, et al. Somatostatin versus octreotide in the treatment of patients with gastrointestinal and pancreatic fistulas. *Can J Gastroenterol*. 2004;18(5):303-306.
36. Allen PJ, Gonen M, Brennan MF, et al. Pasireotide for postoperative pancreatic fistula. *N Engl J Med*. 2014;370(21):2014-2022.

octreotide (Sandostatin), Sandostatin LAR\_PA\_ALL\_MBRx

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.